



Table of Contents

Prologue	7
1. Do not use diagnostic labels initially in front of a patient consulting for emotional distress	9
2. Do not tackle mental health problems without contextualizing first the social health determinants to avoid medicalizing issues with a social basis	11
3. Do not medicalize uncomplicated pain	13
4. Do not minimize symptoms and their impact on mental health of the paediatric and young population	15
5. Do not forget substance consumption in the history of a patient consulting due to a mental health problem	17
6. Do not omit tackling suicidal ideas in people consulting for depressive symptoms	19
7. Do not diagnose a mental health problem without having previously ruled out organic causes	21
8. Do not use psychomedicines exclusively to treat mental health problems	23
9. Do not prolong situations of temporary incapacity for depressive disorders whilst waiting for a full remission of the symptoms	25
10. Do not prescribe benzodiazepines without considering their half-life and the patient's clinical profile beforehand	27
11. Do not prescribe antidepressants systematically for the initial treatment of mild depressive disorder	29
12. If there is no depression do not treat insomnia symptoms with antidepressants that have a sedative profile	31
13. Do not underestimate the cardiovascular risk of the patient treated with antipsychotics	33
14. Do not continue treatments with psychotropic drugs indefinitely without reviewing of the patients' prescription criteria and situation	35
15. Do not deprescribe antidepressants suddenly to avoid discontinuation syndrome	37